arXiv:1607.00895v1 [math.AG] 4 Jul 2016

# ULRICH BUNDLES ON SURFACES WITH $q = p_g = 0$

ARNAUD BEAUVILLE

ABSTRACT. We prove that any surface with $q = p_g = 0$ embedded by a sufficiently large linear system admits a rank 2 Ulrich bundle. In particular every Enriques surface admits a rank 2 Ulrich bundle.

Let $X \subset \mathbb{P}^n$ be a complex projective variety of dimension $d$. A *Ulrich bundle* on $X$ is a vector bundle $E$ on $X$ satisfying $H^\bullet(X, E(-1)) = \ldots = H^\bullet(X, E(-d)) = 0$. This notion was introduced in [ES], where various other characterizations are given; let us just mention that it is equivalent to say that $E$ admits a linear resolution as a $\mathscr{O}_{\mathbb{P}^n}$-module, or that the pushforward of $E$ onto $\mathbb{P}^d$ by a general linear projection is a trivial bundle.

In [ES] the authors ask whether every projective variety admits a Ulrich bundle. The answer is known only in a few cases: hypersurfaces and complete intersections [HUB], del Pezzo surfaces [ES, Corollary 6.5], abelian surfaces [B], sufficiently general K3 surfaces [AFO]. We show here that the Lazarsfeld-Mukai construction produces a rank 2 Ulrich bundle on any surface with $q = p_g = 0$, provided the linear system $|\mathscr{O}_S(1)|$ is large enough :

**Theorem.** *Let $S \subset \mathbb{P}^n$ be a surface with $q = p_g = 0$. Assume that the linear system $|K_S^{-1}(1)|$ contains an irreducible curve. Then $S$ admits a rank 2 Ulrich bundle.*

*In particular, every Enriques surface $S \subset \mathbb{P}^n$ admits a rank 2 Ulrich bundle.*

*Proof* : Let $C$ be an irreducible curve in $|K_S^{-1}(1)|$; we have $\mathscr{O}_S(1)_{|C} = \omega_C$. Let $A$ be a general line bundle of degree $g(C) + 1$ on $C$. Then $h^0(A) = 2$, $h^1(A) = 0$, and $A$ is generated by its global sections. Let $E$ be the kernel of the evaluation map $H^0(C, A) \otimes \mathscr{O}_S \twoheadrightarrow A$. This is a rank 2 vector bundle on $S$. From the exact sequence

$$0 \to E \to H^0(C, A) \otimes \mathscr{O}_S \to A \to 0$$

we get $H^\bullet(S, E) = 0$. Since $\det E = \mathscr{O}_S(-C)$, $E(1)$ is isomorphic to $E^* \otimes K_S$, thus $H^\bullet(S, E(1)) = 0$. Therefore $E(2)$ is a Ulrich bundle for $S$. ∎

*Remarks*.– 1) It might be that every Enriques surface carries actually a Ulrich line bundle. In [BN] the authors conjecture that Enriques surfaces with no $(-2)$-curves admit a Ulrich line bundle, and prove that conjecture in a significant case.

2) Minimal surfaces with Kodaira dimension 0 fall into 4 classes, namely K3, Enriques, abelian and bielliptic surfaces. A bielliptic surface $S$ admits a finite étale covering $\pi : A \to S$ by an abelian surface $A$; if $L$ is a very ample line bundle on $S$ and $E$ a Ulrich bundle for $(A, \pi^* L)$, then $\pi_* E$ is a Ulrich bundle for $(S, L)$. Thus all minimal surfaces with Kodaira dimension 0 admit a Ulrich bundle, except possibly some special K3 surfaces.

3) Let $S \subset \mathbb{P}^n$ be a surface with $q = p_g = 0$, such that $\mathscr{O}_S(1) \cong K_S^r$, with $r \geq 3$. One can show that the linear system $|K_S^{r-1}|$ contains an irreducible curve, hence $S$ admits a rank 2 Ulrich bundle.

## REFERENCES

[AFO] M. Aprodu, G. Farkas, A. Ortega : *Minimal resolutions, Chow forms and Ulrich bundles on $K3$ surfaces*. J. Reine Angew. Math., to appear. Preprint `arXiv:1212.6248`.

[B] A. Beauville : *Ulrich bundles on abelian surfaces*. Proc. Amer. Math. Soc., to appear. Preprint `arXiv:1512.00992`.

[BN] L. Borisov, H. Nuer : *Ulrich Bundles on Enriques Surfaces*. Preprint `arXiv:1606.01459`.

[ES] D. Eisenbud, F.-O. Schreyer : *Resultants and Chow forms via exterior syzygies*. J. Amer. Math. Soc. **16** (2003), no. 3, 537-579.

[HUB] J. Herzog, B. Ulrich, J. Backelin : *Linear maximal Cohen-Macaulay modules over strict complete intersections*. J. Pure Appl. Algebra **71** (1991), no. 2-3, 187-202.

LABORATOIRE J.-A. DIEUDONNÉ, UMR 7351 DU CNRS, UNIVERSITÉ DE NICE, PARC VALROSE, F-06108 NICE CEDEX 2, FRANCE

*E-mail address*: `arnaud.beauville@unice.fr`